



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Short communication

Disinfectant-induced hormesis: An unknown environmental threat of the application of disinfectants to prevent SARS-CoV-2 infection during the COVID-19 pandemic? ☆

Evgenios Agathokleous^{a,*}, Damià Barceló^{b,c}, Ivo Iavicoli^d, Aristidis Tsatsakis^e, Edward J. Calabrese^f

^a Department of Ecology, School of Applied Meteorology, Nanjing University of Information Science and Technology (NUIST), Ningliu Rd. 219, Nanjing, Jiangsu, 210044, China

^b Institute of Environmental Assessment and Water Research, IDAEA-CSIC, C/ Jordi Girona 18-26, 08034, Barcelona, Spain

^c Catalan Institute for Water Research, ICRA-CERCA, Emili Grahit 101, 17003, Girona, Spain

^d Department of Public Health, Section of Occupational Medicine, University of Naples Federico II, Naples, 80131, Italy

^e Laboratory of Toxicology, Medical School, University of Crete, Greece

^f Department of Environmental Health Sciences, Morrill I, N344, University of Massachusetts, Amherst, MA, 01003, USA



ARTICLE INFO

Keywords:

Coronavirus
Disinfectants application
Dose-response relationship
Environmental pollution
Hormesis
Pandemic

ABSTRACT

Massive additional quantities of disinfectants have been applied during the COVID-19 pandemic as infection preventive and control measures. While the application of disinfectants plays a key role in preventing the spread of SARS-CoV-2 infection, the effects of disinfectants applied during the ongoing pandemic on non-target organisms remain unknown. Here we collated evidence from multiple studies showing that chemicals used for major disinfectant products can induce hormesis in various organisms, such as plants, animal cells, and microorganisms, when applied singly or in mixtures, suggesting potential ecological risks at sub-threshold doses that are normally considered safe. Among other effects, sub-threshold doses of disinfectant chemicals can enhance the proliferation and pathogenicity of pathogenic microbes, enhancing the development and spread of drug resistance. We opine that hormesis should be considered when evaluating the effects and risks of such disinfectants, especially since the linear-no-threshold (LNT) and threshold dose-response models cannot identify or predict their effects.

1. Introduction

The world is currently facing the fifth pandemic after the 1918 flu pandemic, which is caused by the novel coronavirus disease 2019 (COVID-19) (Liu et al., 2020; Neagu et al., 2021). The first symptom onset might have occurred on 1 December 2019, and while the press initially called the disease Wuhan pneumonia, whole-genome sequencing revealed that a novel coronavirus was responsible for the disease (Liu et al., 2020). The World Health Organization (WHO) named the virus ‘2019 novel coronavirus’ (2019-nCoV) in mid January 2020, following by its official declaration as ‘coronavirus disease 2019’ (COVID-19) one month later (Liu et al., 2020). Some weeks later, the Coronaviridae Study Group (CSG) of the International Committee on

Taxonomy of Viruses (ICTV) finally recognized the virus as “forming a sister clade to the prototype human and bat severe acute respiratory syndrome coronaviruses (SARS-CoVs) of the species *Severe acute respiratory syndrome-related coronavirus*”, designated as ‘SARS-CoV-2’ (CSG-ICTV, 2020). As of 29 October 2021, 246,245,186 cases of infection, 223,134,984 recoveries, and 4,995,890 deaths have been recorded across the globe (<https://www.worldometers.info/coronavirus/>; last updated: 29 October 2021, 01:21 GMT).

To prevent further spreading of SARS-CoV-2, governments all over the world imposed mobility restriction measures of various degrees of severity, including local, regional, or national lockdowns (Benchrif et al., 2021; Calina et al., 2021; Deroubaix et al., 2021; Kim et al., 2021; Marinello et al., 2021), and mandated the application of public

☆ This paper has been recommended for acceptance by Christian Sonne.

* Corresponding author.

E-mail address: evgenios@nuist.edu.cn (E. Agathokleous).

health-based measures such as wearing masks and sanitizing public spaces (Dadras et al., 2021; Subpiramanyam, 2021). Strategies to halt the infection cycle have included the disinfection of objects and surfaces (Barcelo, 2020; Khan and Yadav, 2020; Kwok et al., 2021). As a direct consequence of this strategy and its widespread implementation, disinfectants have been massively applied across the globe, and the extent of this practice is considerable. For example, disinfectants should be applied multiple times per day in contaminated sites, with each treatment lasting enough time per application (e.g. at least 30 min for 2000 mg/L chlorine-containing disinfectants) (Barcelo, 2020), while China had dispensed 2000–5000 tons of disinfectants in Wuhan alone (Zhang et al., 2020). The effectiveness of the disinfectants to inactivate SARS-CoV-2 depends on various factors, however, many studies show that such chemical products are efficient in inactivating the virus if applied properly (Dhama et al., 2021; Sharafi et al., 2021; Shimabukuro et al., 2020).

While the effectiveness to reduce significantly the spread of SARS-CoV-2 is an area of scientific research, there are potential side effects on the environment, affecting various creatures such as plants, insects, and wild animals (Dhama et al., 2021; Ghafoor et al., 2021). Several studies examine in detail the potential impacts of the application of such disinfectants to the environment (Ankit et al., 2021; Bonin et al., 2020; Chen et al., 2021; Dhama et al., 2021; Nabi et al., 2020; Zhang et al., 2020). There is, however, no study evaluating the effects of such chemicals within the context of dose-response relationship and whether such chemicals can induce hormesis in various living organisms, despite the high importance of this for risk assessment considerations (Agathokleous and Calabrese, 2020a). Here, we aimed at examining the published literature for evidence supporting the occurrence of hormesis as a result of exposure to compounds used in disinfectants applied to surfaces, as well as ethanol-based compounds used as preventive measures against SARS-CoV-2 infection. Because disinfectants applied to surfaces include many chemicals other than ethanol that are longer-lived and thus expected to have wider environmental consequences, we focused on the potential implications of hormesis induced by surface disinfectants. The herein analysis suggests that the massive application of disinfectants for containing SARS-CoV-2 may be a double-edged sword, inhibiting/preventing the virus but also imposing some potentially significant but non-apparent costs or risks by affecting other non-target organisms in a dose-dependent manner, and finally promoting traits of drug resistance.

2. Presentation of the analysis and concerns

The massive release of SARS-CoV-2 disinfectants into the waste (water) systems and run-off continuously since early 2020 is alarming in terms of potential effects on non-target species (other than SARS-CoV-2) (Subpiramanyam, 2021). Collectively, the WHO considered the application of chlorine-based chemicals for disinfecting health-care settings and other environmental surfaces (WHO, 2020). Moreover, the US Environmental Protection Agency (EPA) is continuously updating the disinfectants list for application onto surfaces (acknowledged also by WHO (2020)), including various chemicals, such as H₂O₂ alone or with peroxyacetic acid (peracetic acid), dodecylbenzenesulfonic acid alone or with L-lactic acid, quaternary ammonium alone or with isopropanol (isopropyl alcohol), and sodium carbonate peroxyhydrate with tetraacetyl ethylenediamine (<https://cfpub.epa.gov/wizards/disinfectants>; Accessed on 3 August 2021). A large proportion of the EPA's recommended disinfectant products for application against SARS-CoV-2 is based on quaternary ammonium compounds (QACs) (Hora et al., 2020). Although QACs were extensively applied and detected in sediments, surface waters, and wastewater before the current pandemic, their use

has increased during the SARS-CoV-2 pandemic, which is of more concern due to their role into antibiotic resistance (Hora et al., 2020; Jiang et al., 2022; Zhang et al., 2015).¹ Additional disinfectants that can be applied to halt SARS-CoV spread are chemicals such as ethanol (Dhama et al., 2021; Hirose et al., 2020; Ijaz et al., 2021; Kwok et al., 2021; Rabenau et al., 2005; Xiling et al., 2021).

Massive application of disinfectants against COVID-19, however, can have various negative effects on humans and the environment, poses potential environmental/ecological risks, especially in urban ecosystems, and may undermine the concept of 'One Health' (Nabi et al., 2020). For example, disinfectants can affect the mucosal lining, thus leading to inflammation, irritation, swelling, and ulceration of the respiratory tract (Dhama et al., 2021). They can also cause skin dryness and heighten the risk of developing asthma, chronic obstructive pulmonary disease, and impaired brain development, and infertility in children (Dhama et al., 2021). In addition to the effects on humans, a large discharge of disinfectants into lakes, rivers, sewage, due to activities including the cleaning of external floors, markets, and streets (Subpiramanyam, 2021), and disinfection of wastewater derived from heavy activities, such as healthcare facilities, hotels, and factories, can affect living organisms (Dhama et al., 2021). For example, chlorine-based disinfectants may (i) oxidize proteins and destruct cell walls of aquatic wildlife and plants, (ii) generate byproducts, e.g. haloacetic acids and trihalomethanes, which may be highly toxic to aquatic flora and fauna, (iii) bond with other contaminants and transform into harmful compounds, and (iv) potentially affecting the activity of pollutant-removing microorganisms in wastewater treatment plants (Dhama et al., 2021; Sedlak and Von Gunten, 2011; Subpiramanyam, 2021; Zhang et al., 2020). In the worst case scenario, animals may even die due to overload of disinfectants, as it has been reported after application of massive amounts of disinfectants to a coronavirus epicenter (Nabi et al., 2020). Therefore, the ongoing massive application of disinfectants may threaten numerous creatures in terrestrial and aquatic systems (Zhang et al., 2020), urging for evaluations of dose-response relationships in a plethora of living organisms exposed to disinfectants.

Although the effects of such disinfectants to non-target organisms remain unknown, there is evidence that such chemicals can induce hormesis with significant responses at sub-threshold doses, i.e. at doses smaller than the traditional toxicological threshold (Fig. 1). Examples of such chemicals shown to induce hormesis include chlorine-based chemicals (Zhang et al., 2008), H₂O₂ (Huang et al., 2019; Ludovico and Burhans, 2014; Semchyshyn, 2014; Semchyshyn and Valishkevych, 2016), various QACs (Hrubec et al., 2021; Li et al., 2019; Mo et al., 2020), and ethanol (Calabrese and Baldwin, 2003; Kar et al., 2021; Semchyshyn, 2014). These chemicals are used for both hand and surface sanitizer products and are among the most widely applied disinfectants during the current pandemic (Dhama et al., 2021; Jing et al., 2020; World Health Organization (WHO), 2020). Importantly, hormesis was induced by not only single chemicals but also mixtures, specifically in studies with QACs, which can display also additive or synergistic effects (Mo et al., 2020); hormetic effects of complex mixtures of chemicals have also been reported by numerous other studies (Docea et al., 2019; Tsatsakis et al., 2019). Extensive experimentation with yeast subjected to a series of dilutions of numerous disinfectants revealed that most of the disinfectants induced toxicity followed by an overcompensation stimulation (Branham, 1929). The occurrence of overcompensation stimulation within a hormetic framework is very common, and highly generalizable (Calabrese, 2001). These findings suggest that both direct stimulation and overcompensation can occur in the framework of disinfectant-induced hormesis, which would make the concern greater due to the complexity to identify and predict such effects. Newer studies incorporating a time component would be needed to examine

¹ See also the information provided by the US EPA at <https://www.epa.gov/coronavirus/disinfectant-use-and-coronavirus-covid-19>.




Fig. 1. A: A schematic representation of hormetic dose-response relationship. Similar hormetic-like responses were found in various organisms exposed to different types of disinfectant-containing chemicals. B: An example of hormetic dose-response relationship. Cell viability assay was evaluated with the MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) tetrazolium reduction assay in a mouse embryonic fibroblast cell line. The figure has been created based on data reported by Kar et al. (2021), and asterisk above a mean indicates significant difference from the control based on the original authors' statistical analyses. The ethanol concentrations were transformed for presentation purposes. The original concentrations were 0, 0.005, 0.01, 0.05, 0.1, 0.5, 1, 5, and 10 v/v % for 48 h. Data were extracted from the relevant figure of the published article using Adobe Photoshop CS4 Extended v.11 (Adobe Systems Incorporated, CA, USA).

disinfectant-induced time-dependent hormesis in different organisms and endpoints.

The preceding collated evidence indicates the induction of hormesis by various disinfectants. Hormesis has also been widely observed in various living organisms exposed to numerous other pollutants and contaminants of emerging concern of aquatic and terrestrial systems, including active pharmaceutical ingredients, formaldehyde, fungicides, heavy metals and other elements, herbicides, hydrocarbons, microplastics, nanomaterials, and ozone (Agathokleous et al., 2021c; Agathokleous and Calabrese, 2020b, 2021; Belz and Duke, 2017; Calabrese and Agathokleous, 2021; Carvalho et al., 2020; Erofeeva, 2021a, 2021b; Iavicoli et al., 2021; Shahid et al., 2020). The induction of hormesis by so many contaminants with considerably differed molecular structure suggests the possibility that any chemical agent may induce hormesis when applied at the right doses. Nevertheless, more studies are needed to (i) examine whether all the disinfectants that are currently available and applied induce hormesis in various creatures, (ii) reveal what the underlying mechanisms are, and (iii) understand whether the underlying mechanisms and quantitative traits of hormetic responses differ among disinfectants.

Hormesis was induced in various organisms, such as freshwater luminescent bacteria (*Vibrio qinghaiensis* sp. Q67) (Mo et al., 2020), hydroponically-cultivated higher plants (*Triticum aestivum* L.) (Li et al., 2019), human neuroblastoma cell line (Huang et al., 2019), humans (associational) (Hrubec et al., 2021), mouse embryonic fibroblast cells (Kar et al., 2021), mouse macrophages (Zhang et al., 2008), and yeast (*Saccharomyces cerevisiae* Meyen ex E.C. Hansen) (Semchyshyn, 2014; Semchyshyn and Valishkevych, 2016). These recent findings extend older findings of ethanol- and other disinfectant-induced hormesis in

different kinds of cells and experimental models (Branham, 1929; Calabrese and Baldwin, 2003), and the magnitude of the sub-threshold effects is commonly within two-fold of the control response/status, in agreement with the broad hormesis literature for numerous stresses, organisms, and endpoints (Agathokleous et al., 2020; Calabrese et al., 2019; Calabrese and Agathokleous, 2021; Sun et al., 2021). Hormetic responses have been reported for a plethora of endpoints, such as cell viability (Zhang et al., 2008) and proliferation [MTT activity (Kar et al., 2021)], cyanobacteria luciferase (Mo et al., 2020), plant shoot and root biomass (Li et al., 2019), human health-related biomarkers (Hrubec et al., 2021), and levels of reduced glutathione (GSH) and gene expression of GCLC and NQO-1 (Zhang et al., 2008). H₂O₂ also promoted PAC1-R promoter activity in a dose-dependent fashion, which was inhibited by the transcription factor specificity protein 1 (SP1) inhibitor mithramycin A (Huang et al., 2019), while yeast colony growth positively correlated with the glutathione reductase activity (Semchyshyn, 2014; Semchyshyn and Valishkevych, 2016). Not only can cells show adaptability to sub-threshold doses of ethanol (Kar et al., 2021), but also H₂O₂ applied at sub-threshold doses could lead to cross-resistance of budding yeast (*S. cerevisiae*) to different stresses, a phenomenon where the regulatory protein Yap1 plays an important role (Semchyshyn, 2014). Although hormetic effects depend on the carbohydrate (energy source) in growth medium (Semchyshyn and Valishkevych, 2016), these findings suggest that low doses of disinfectants in the environment may change the sensitivity of non-target organisms to different environmental stresses.

These results also suggest that sub-threshold doses of disinfectants can have various biologically positive effects on non-target organisms, like many other contaminants (Agathokleous et al., 2021c, 2020; Calabrese and Agathokleous, 2021; Iavicoli et al., 2021); however, with the potential to translate to ecologically/environmentally negative outcomes. For example, while they enhance plant biomass, they can stimulate emerging fungal pathogens (Pérez-Torrado and Querol, 2016) and perplex their control in specific environmental settings, especially if such sub-threshold responses are linked with mechanisms involved in the development of resistance (Iavicoli et al., 2021). Bacterial populations exhibit high heterogeneity and are composed of cells of various ages and physiological states, while antibiotics effects on bacteria might be defined by how many molecules effectively interact with individual cells (Baquero and Levin, 2020). Of increased concern is that there is ample evidence indicating that QACs can (i) lead to proliferation of pathogenic, multidrug-resistant bacteria and cross-resistance of bacteria to antibiotics, (ii) emergence of microflora enriched with bacterial strains presenting resistance to clinically important antimicrobial agents, and (iii) promotion of the fixation of novel genetic elements, which assist in spreading resistance genes (Akimitsu et al., 1999; Braoudaki and Hilton, 2004; Buffet-Bataillon et al., 2012; Chen et al., 2021; Hora et al., 2020; Pereira and Tagkopoulos, 2019; Tezel and Pavlostathis, 2011; Zhang et al., 2015). These may add to the existing issue of the increase in multidrug-resistant organisms (Exner et al., 2020; see also Berendonk et al., 2015). Even if some disinfectants such as ethanol may not lead to environmental consequences, it is still possible that bacterial resistance may develop in the bacterial flora associated with the skin of humans and other animals (Chan, 1999; Chen et al., 2021; Ghafoor et al., 2021; Schwarz et al., 2017), a hypothesis that remains to be studied.

It should also be mentioned, however, that different disinfectants may have distinct extent of spatiotemporal impact of the longer-lived compounds, such as QACs, versus the shorter-lived chemicals, such as H₂O₂ and ethanol (Dhama et al., 2021). The reduced longevity of H₂O₂ and ethanol, e.g. due to its sensitivity to light or its high vapor pressure, may be expected to limit the consequences to local environments and skin (hence people), whilst the longer-lived compounds may be expected to have wider-reaching environmental effects (ecosystems). Importantly, there is a wealth of studies illustrating that exposure of various creatures to low doses, which are smaller than the traditional

toxicological threshold, can have long-lasting effects, even transmitted transgenerationally and observed even more than ten generations later (Agathokleous et al., 2021b; Agathokleous and Calabrese, 2020b; Calabrese and Agathokleous, 2020). These suggest that effects of massively applied disinfectants might persist for a long time in the environment, indicating a need for new-generation (eco)toxicological studies that would be directed to reveal potential transgenerational effects of sub-toxic doses of disinfectants. Therefore, hormesis should be considered in risk assessment because the traditional and biologically irrelevant threshold and linear-no-threshold (LNT) dose response models cannot capture effects that occur in the hormetic (low-dose) zone (Agathokleous et al., 2021a, 2019; Agathokleous and Calabrese, 2020a; Tsatsakis, 2021). No consideration of hormetic responses may lead to considerably incorrect estimates of points of departure, while the complexity of disinfectant and other chemical mixtures and the non-linear hormetic responses would make the derivation of reliable permissible levels challenging for risk assessment-based regulation (Agathokleous et al., 2021d; Tsatsakis, 2021).

Disinfectants are also widely used for drinking water treatment (Humans, 1991; Pichel et al., 2019; Tsitsifli and Kanakoudis, 2018), especially in the US, South Europe, and China. Hence, in several countries chlorination takes place in the drinking water plant and at the effluent to ensure that the drinking water network is charged with enough chlorine when reaching the public. The recent massive documentation of hormetic responses induced by disinfectants and other chemicals suggests the possibility that sub-toxic doses of such chemicals in the tap water might have unknown effects that traditional dose-response models fail to identify. Therefore, new generation studies are needed to evaluate for potential effects of disinfectants added into the tap water at doses that are smaller than the traditional toxicological threshold. A complementary tool for assessing the whole treatment process is by using a group of toxicity tests of different end-points, thus controlling the toxicity or effects of a plethora of pollutants present in water (Barceló et al., 2020).

3. Conclusions

The use of disinfectants has increased amid the ongoing pandemic and is expected to remain elevated (Hora et al., 2020), raising concerns regarding the fate of such chemicals into the environment. In this paper, we collated considerable evidence for the occurrence of hormesis in various organisms exposed to single or combined chemicals used for the production of disinfectants that are widely used during the current pandemic. Such significant sub-threshold responses are of profound importance to be acknowledged, identified, and predicted to avoid potential negative consequences to the environment, ecological processes, and structure and function of communities and ecosystems.

Real-world samples, however, contain a variety of chemicals, not only one specific group of chemicals, and the study of non-linear dose-response relationships induced by complex mixtures is challenging and practically difficult (Docea et al., 2021; Kumari and Kumar, 2020). Inadequate information about the combined exposure to such chemical mixtures also hampers risk assessment under a real-life risk simulation (RLRS) approach (Agathokleous et al., 2019; Hernández et al., 2020). Although this analysis shows hormesis induced by various chemical mixtures, in agreement with the wider literature documenting hormesis induced by diverse chemical mixtures (Agathokleous et al., 2020), new studies are needed to investigate non-linear dose-response relationships induced by diverse mixtures containing disinfectants combined with various other types of chemicals.

Amongst other issues, hormetic effects of disinfectants may stimulate potentially pathogenic bacteria and facilitate the development and spread of drug resistance, implications for risk assessment that can be tackled only if hormetic responses are taken into account. The increasing use of antibiotics around the world (Aslam et al., 2018; Zaman et al., 2017), perplexes the antibiotic removal by the wastewater treatment

plants (WWTPs) as well as the antibiotic resistance genes. The aquatic environment already contains traces of antibiotic resistance genes, especially when large hospitals discharge in WWTPs (Kumar et al., 2021; Rodriguez-Mozaz et al., 2015). Hence, assuming that the application of SARS-CoV-2 disinfectants during the COVID-19 pandemic may worsen the situation by adding to the antibiotic resistance issue, advanced WWTP might be needed in addition to changing disinfection formulations.

In conclusion, the present exposition indicates a dilemma of the extensive application of disinfectants against specific human pathogens due to their ability to induce hormesis in various non-target organisms. It also highlights the need for the discovery of alternative products with equal or higher disinfecting ability but with less potential side-effects in order to be prepared when such a massive application of disinfectants is unavoidable.

Credit author statement

Evgenios Agathokleous: Writing- Original draft preparation, Visualization. Damià Barceló: Writing- Reviewing and Editing. Ivo Iavicoli: Writing- Reviewing and Editing. Aristides Tsatsakis: Writing- Reviewing and Editing. Edward J Calabrese: Writing- Reviewing and Editing.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. E.A. acknowledges multi-year support from The Startup Foundation for Introducing Talent of Nanjing University of Information Science & Technology (NUIST), Nanjing, China (Grant No. 003080). E.J.C. acknowledges longtime support from the US Air Force (Grant No. AFOSR FA9550-13-1-0047) and ExxonMobil Foundation (Grant No. S1820000000256). The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing policies or endorsement, either expressed or implied. Sponsors had no involvement in study design, collection, analysis, interpretation, writing and decision to and where to submit for publication consideration.

Author contributions

E.A. drafted the manuscript, had a leading role, served as the hub of communication among the authors, and supervised the production of the manuscript. D.B., I.I., A.T., and E.J.C. reviewed the manuscript and contributed intellectual input. All authors approved the final version for publication.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Agathokleous, E., Anav, A., Aramineni, V., De Marco, A., Domingos, M., Kitao, M., Koike, T., Manning, W.J., Paoletti, E., Saitanis, C.J., Sicard, P., Vitale, M., Wang, W., Calabrese, E.J., 2019. Commentary: EPA's proposed expansion of dose-response analysis is a positive step towards improving its ecological risk assessment. *Environ. Pollut.* 246, 566–570. <https://doi.org/10.1016/j.envpol.2018.12.046>.
- Agathokleous, E., Barceló, D., Calabrese, E.J., 2021a. US EPA: is there room to open a new window for evaluating potential sub-threshold effects and ecological risks? *Environ. Pollut.* 284, 117372. <https://doi.org/10.1016/j.envpol.2021.117372>.
- Agathokleous, E., Brown, P.H., Calabrese, E.J., 2021b. A gift from parent to offspring: transgenerational hormesis. *Trends Plant Sci.* 26, 1098–1100. <https://doi.org/10.1016/j.tplants.2021.08.006>.
- Agathokleous, E., Calabrese, E.J., 2021. Fungicide-induced hormesis in phytopathogenic fungi: a critical determinant of successful agriculture and environmental

- sustainability. *J. Agric. Food Chem.* 69, 4561–4563. <https://doi.org/10.1021/acs.jafc.1c01824>.
- Agathokleous, E., Calabrese, E.J., 2020a. Environmental toxicology and ecotoxicology: how clean is clean? Rethinking dose-response analysis. *Sci. Total Environ.* 746, 138769. <https://doi.org/10.1016/j.scitotenv.2020.138769>.
- Agathokleous, E., Calabrese, E.J., 2020b. A global environmental health perspective and optimization of stress. *Sci. Total Environ.* 704, 135263. <https://doi.org/10.1016/j.scitotenv.2019.135263>.
- Agathokleous, E., Iavicoli, I., Barceló, D., Calabrese, E.J., 2021c. Micro/nanoplastics effects on organisms: a review focusing on 'dose. *J. Hazard Mater.* 417, 126084. <https://doi.org/10.1016/j.jhazmat.2021.126084>.
- Agathokleous, E., Kitao, M., Calabrese, E.J., 2020. Hormesis: highly generalizable and beyond laboratory. *Trends Plant Sci.* 25, 1076–1086. <https://doi.org/10.1016/j.tplants.2020.05.006>.
- Agathokleous, E., Saitanis, C., Markouizou, A., 2021d. Hormesis shifts the no-observed-adverse-effect level (NOAEL). *Dose-Response* 19. <https://doi.org/10.1177/15593258211001667>, 15593258211001668.
- Akimitsu, N., Hamamoto, H., Inoue, R., Shoji, M., Akamine, A., Takemori, K., Hamasaki, N., Sekimizu, K., 1999. Increase in resistance of methicillin-resistant *Staphylococcus aureus* to beta-lactams caused by mutations conferring resistance to benzalkonium chloride, a disinfectant widely used in hospitals. *Antimicrob. Agents Chemother.* 43, 3042–3043. <https://doi.org/10.1128/aac.43.12.3042>.
- Ankit Kumar, A., Jain, V., Deovanshi, A., Lepcha, A., Das, C., Baudhdh, K., Srivastava, S., 2021. Environmental impact of COVID-19 pandemic: more negatives than positives. *Environ. Sustain.* 1 <https://doi.org/10.1007/S42398-021-00159-9>.
- Aslam, B., Wang, W., Arshad, M.I., Khurshid, M., Muzammil, S., Rasool, M.H., Nisar, M. A., Alvi, R.F., Aslam, M.A., Qamar, M.U., Salamat, M.K.F., Baloch, Z., 2018. Antibiotic resistance: a roundup of a global crisis. *Infect. Drug Resist.* 11, 1645. <https://doi.org/10.2147/idr.s173867>.
- Baquero, F., Levin, B.R., 2020. Proximate and ultimate causes of the bactericidal action of antibiotics. *Nat. Rev. Microbiol.* 19, 123–132. <https://doi.org/10.1038/s41579-020-00443-1>, 2020.
- Barceló, D., 2020. An environmental and health perspective for COVID-19 outbreak: meteorology and air quality influence, sewage epidemiology indicator, hospitals disinfection, drug therapies and recommendations. *J. Environ. Chem. Eng.* 8, 104006. <https://doi.org/10.1016/j.jece.2020.104006>.
- Barceló, D., Zonja, B., Ginebreda, A., 2020. Toxicity tests in wastewater and drinking water treatment processes: a complementary assessment tool to be on your radar. *J. Environ. Chem. Eng.* 8, 104262. <https://doi.org/10.1016/j.jece.2020.104262>.
- Belz, R.G., Duke, Stephen O., 2017. Herbicide-Mediated hormesis. In: Duke, S.O., Kudsk, P., Solomon, K. (Eds.), *Pesticide Dose: Effects on the Environment and Target and Non-target Organisms*. American Chemical Society, Washington, pp. 135–148. <https://doi.org/10.1021/bk-2017-1249.ch010>.
- Benchrif, A., Wheida, A., Tahri, M., Shubbar, R.M., Biswas, B., 2021. Air quality during three covid-19 lockdown phases: AQI, PM_{2.5} and NO₂ assessment in cities with more than 1 million inhabitants. *Sustain. Cities Soc.* 74, 103170. <https://doi.org/10.1016/j.scs.2021.103170>.
- Berendonk, T.U., Manaia, C.M., Merlín, C., Fatta-Kassinos, D., Cytryn, E., Walsh, F., Bürgmann, H., Sørum, H., Norström, M., Pons, M.N., Kreuzinger, N., Huovinen, P., Stefani, S., Schwartz, T., Kisan, V., Baquero, F., Martinez, J.L., 2015. Tackling antibiotic resistance: the environmental framework. *Nat. Rev. Microbiol.* 13, 310–317. <https://doi.org/10.1038/nrmicro3439>.
- Bonin, L., Vitry, V., Olivier, M.-G., Bertolucci-Coeelho, L., 2020. Covid-19: effect of disinfection on corrosion of surfaces. *Corrosion Eng. Sci. Technol.* 55, 693–695. <https://doi.org/10.1080/1478422x.2020.1777022>.
- Branham, S.E., 1929. The effects of certain chemical compounds upon the course of gas production by baker's yeast. *J. Bacteriol.* 18, 247–264. <https://doi.org/10.1128/jb.18.4.247-264.1929>.
- Braoudaki, M., Hilton, A.C., 2004. Adaptive resistance to biocides in *Salmonella enterica* and *Escherichia coli* O157 and cross-resistance to antimicrobial agents. *J. Clin. Microbiol.* 42, 73–78. <https://doi.org/10.1128/jcm.42.1.73-78.2004>.
- Buffet-Bataillon, S., Tattevin, P., Bonnaure-Mallet, M., Jolivet-Gougeon, A., 2012. Emergence of resistance to antibacterial agents: the role of quaternary ammonium compounds—a critical review. *Int. J. Antimicrob. Agents* 39, 381–389. <https://doi.org/10.1016/j.ijantimicag.2012.01.011>.
- Calabrese, E.J., 2001. Overcompensation stimulation: a mechanism for hormetic effects. *Crit. Rev. Toxicol.* 31, 425–470. <https://doi.org/10.1080/20014091111749>.
- Calabrese, E.J., Agathokleous, E., 2021. Accumulator plants and hormesis. *Environ. Pollut.* 274, 116526. <https://doi.org/10.1016/j.envpol.2021.116526>.
- Calabrese, E.J., Agathokleous, E., 2020. Theodosius Dobzhansky's view on biology and evolution v.2.0: "Nothing in biology makes sense except in light of evolution and evolution's dependence on hormesis-mediated acquired resilience that optimizes biological performance and numerous diverse short and longer term protective strategies". *Environ. Res.* 186, 109559. <https://doi.org/10.1016/j.envres.2020.109559>.
- Calabrese, E.J., Agathokleous, E., Kozumbo, W.J., Stanek, E.J., Leonard, D., 2019. Estimating the range of the maximum hormetic stimulatory response. *Environ. Res.* 170, 337–343. <https://doi.org/10.1016/j.envres.2018.12.020>.
- Calabrese, E.J., Baldwin, L.A., 2003. Ethanol and hormesis. *Crit. Rev. Toxicol.* 33, 407–424. <https://doi.org/10.1080/103611043>.
- Calina, D., Hartung, T., Mardare, I., Mitroi, M., Poulas, K., Tsatsakis, A., Rogoveanu, I., Docea, A.O., 2021. COVID-19 pandemic and alcohol consumption: impacts and interconnections. *Toxicol. Reports* 8, 529–535. <https://doi.org/10.1016/j.toxrep.2021.03.005>.
- Carvalho, M.E.A., Castro, P.R.C., Azevedo, R.A., 2020. Hormesis in plants under Cd exposure: from toxic to beneficial element? *J. Hazard Mater.* 384, 121434. <https://doi.org/10.1016/j.jhazmat.2019.121434>.
- Chan, J.C., 1999. Combating bacterial resistance in skin and skin-structure infection: importance of beta-lactamase inhibition. *Am. J. Therapeut.* 6, 13–18. <https://doi.org/10.1097/00045391-199901000-00003>.
- Chen, B., Han, J., Dai, H., Jia, P., 2021. Biocide-tolerance and antibiotic-resistance in community environments and risk of direct transfers to humans: unintended consequences of community-wide surface disinfecting during COVID-19? *Environ. Pollut.* 283, 117074. <https://doi.org/10.1016/j.envpol.2021.117074>.
- CSG-ICTV, 2020. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat. Microbiol.* 5, 536. <https://doi.org/10.1038/S41564-020-0695-z>.
- Dadras, O., Alinaghi, S.A.S., Karimi, A., MohsseniPour, M., Barzegary, A., Vahedi, F., Pashaei, Z., Mirzapour, P., Fakhouri, A., Zargari, G., Saedi, S., Mojdaganlou, H., Badri, H., Qaderi, K., Behzad, F., Mehraeen, E., 2021. Effects of COVID-19 prevention procedures on other common infections: a systematic review. *Eur. J. Med. Res.* 26, 67. <https://doi.org/10.1186/s40001-021-00539-1>, 2021.
- Deroubaix, A., Brasseur, G., Gaubert, B., Labuhn, I., Menut, L., Siour, G., Tuccella, P., 2021. Response of surface ozone concentration to emission reduction and meteorology during the COVID-19 lockdown in Europe. *Meteorol. Appl.* 28, e1990. <https://doi.org/10.1002/met.1990>.
- Dhama, K., Patel, S.K., Kumar, R., Masand, R., Rana, J., Yatoo, M.I., Tiwari, R., Sharun, K., Mohapatra, R.K., Natesan, S., Dhawan, M., Ahmad, T., Emran, T. Bin, Malik, Y.S., Harapan, H., 2021. The role of disinfectants and sanitizers during COVID-19 pandemic: advantages and deleterious effects on humans and the environment. *Environ. Sci. Pollut. Res. Int.* 28, 34211–34228. <https://doi.org/10.1007/S11356-021-14429-w>.
- Docea, A.O., Calina, D., Tsatsakis, A.M., 2021. Chapter 1 - mixture toxicity evaluation in modern toxicology. In: Tsatsakis, A.M. (Ed.), *Toxicological Risk Assessment and Multi-System Health Impacts from Exposure*. Academic Press, pp. 3–12. <https://doi.org/10.1016/b978-0-323-85215-9.00044-1>.
- Docea, A.O., Goumenou, M., Calina, D., Arsene, A.L., Dragoi, C.M., Gofita, E., Pisoschi, C. G., Zlatian, O., Stivaktakis, P.D., Nikolouzakis, T.K., Kalogeraki, A., Izotov, B.N., Galateanu, B., Hudita, A., Calabrese, E.J., Tsatsakis, A., 2019. Adverse and hormetic effects in rats exposed for 12 months to low dose mixture of 13 chemicals: RLRs part III. *Toxicol. Lett.* 310, 70–91. <https://doi.org/10.1016/j.toxlet.2019.04.005>.
- Erofeeva, E.A., 2021a. Plant hormesis and Shelford's tolerance law curve. *J. For. Res.* 32, 1789–1802. <https://doi.org/10.1007/s11676-021-01312-0>.
- Erofeeva, E.A., 2021b. Environmental hormesis of non-specific and specific adaptive mechanisms in plants. *Sci. Total Environ.* 804, 150059. <https://doi.org/10.1016/j.scitotenv.2021.150059>.
- Exner, M., Bhattacharya, S., Gebel, J., Goroncy-Bermes, P., Hartemann, P., Heeg, P., Ilschner, C., Kramer, A., Ling, M.L., Merckens, W., Oltmanns, P., Pitten, F., Rotter, M., Schmitzhausen, R.M., Sonntag, H.-G., Steinhauer, K., Trautmann, M., 2020. Chemical disinfection in healthcare settings: critical aspects for the development of global strategies. *GMS Hyg. Infect. Control* 15, Doc36. <https://doi.org/10.3205/dgkh000371>.
- Ghafoor, D., Khan, Z., Khan, A., Ualiyeva, D., Zaman, N., 2021. Excessive use of disinfectants against COVID-19 posing a potential threat to living beings. *Curr. Res. Toxicol.* 2, 159–168. <https://doi.org/10.1016/j.crtox.2021.02.008>.
- Hernández, A.F., Docea, A.O., Goumenou, M., Sarigiannis, D., Aschner, M., Tsatsakis, A., 2020. Application of novel technologies and mechanistic data for risk assessment under the real-life risk simulation (RLRS) approach. *Food Chem. Toxicol.* 137, 111123. <https://doi.org/10.1016/j.fct.2020.111123>.
- Hirose, R., Ikegaya, H., Naito, Y., Watanabe, N., Yoshida, T., Bandou, R., Daidoji, T., Itoh, Y., Nakaya, T., 2020. Survival of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and Influenza virus on human skin: importance of hand hygiene in coronavirus disease 2019 (COVID-19). *Clin. Infect. Dis.* ciae1517. <https://doi.org/10.1093/cid/ciae1517>.
- Hora, P.I., Pati, S.G., McNamara, P.J., Arnold, W.A., 2020. Increased use of quaternary ammonium compounds during the SARS-CoV-2 pandemic and beyond: consideration of environmental implications. *Environ. Sci. Technol. Lett.* 7, 622–631. <https://doi.org/10.1021/acs.estlett.0c00437>.
- Hrubec, T.C., Seguin, R.P., Xu, L., Cortopassi, G.A., Datta, S., Hanlon, A.L., Lozano, A.J., McDonald, V.A., Healy, C.A., Anderson, T.C., Musse, N.A., Williams, R.T., 2021. Altered toxicological endpoints in humans from common quaternary ammonium compound disinfectant exposure. *Toxicol. Reports* 8, 646–656. <https://doi.org/10.1016/j.toxrep.2021.03.006>.
- Huang, X., Li, J., Song, S., Wang, L., Lin, Z., Ouyang, Z., Yu, R., 2019. Hormesis effect of hydrogen peroxide on the promoter activity of neuropeptide receptor PAC1-R. *J. Food Biochem.* 43, e12877. <https://doi.org/10.1111/jfbc.12877>.
- Humans, I.W.G., on the E. of C.R. to, 1991. Chlorinated drinking-water; chlorination by-products; some other halogenated compounds; cobalt and cobalt compound. *IARC Monogr. Eval. Carcinog. Risks Hum.* 52, 45–141.
- Iavicoli, I., Fontana, L., Agathokleous, E., Santocono, C., Russo, F., Vetrani, I., Fedele, M., Calabrese, E.J., 2021. Hormetic dose responses induced by antibiotics in bacteria: a phantom menace to be thoroughly evaluated to address the environmental risk and tackle the antibiotic resistance phenomenon. *Sci. Total Environ.* 798, 149255. <https://doi.org/10.1016/j.scitotenv.2021.149255>.
- Ijaz, M.K., Nims, R.W., Zhou, S.S., Whitehead, K., Srinivasan, V., Kapes, T., Fanuel, S., Epstein, J.H., Daszak, P., Rubino, J.R., McKinney, J., 2021. Microbicidal actives with virucidal efficacy against SARS-CoV-2 and other beta- and alpha-coronaviruses and implications for future emerging coronaviruses and other enveloped viruses. *Sci. Rep.* 11, 5626. <https://doi.org/10.1038/s41598-021-84842-1>.

- Jiang, Q., Feng, M., Ye, C., Yu, X., 2022. Effects and relevant mechanisms of non-antibiotic factors on the horizontal transfer of antibiotic resistance genes in water environments: A review. *Sci. Total Environ.* 806, 150568 <https://doi.org/10.1016/j.scitotenv.2021.150568>.
- Jing, J.L.J., Yi, T.P., Bose, R.J.C., McCarthy, J.R., Tharmalingam, N., Madheswaran, T., 2020. Hand sanitizers: a review on formulation aspects, adverse effects, and regulations. *Int. J. Environ. Res. Publ. Health* 17, 3326. <https://doi.org/10.3390/ijerph17093326>.
- Kar, N., Gupta, D., Bellare, J., 2021. Ethanol affects fibroblast behavior differentially at low and high doses: a comprehensive, dose-response evaluation. *Toxicol. Reports* 8, 1054–1066. <https://doi.org/10.1016/j.toxrep.2021.05.007>.
- Khan, M.H., Yadav, H., 2020. Sanitization during and after COVID-19 pandemic: a short review. *Trans. Indian Natl. Acad. Eng.* 5, 1. <https://doi.org/10.1007/s41403-020-00177-9>.
- Kim, Y.-K., Cho, J.-H., Kim, H.-S., 2021. Analysis of concentration variations of long-range transport PM10, NO₂, and O₃ due to COVID-19 shutdown in East Asia in 2020. *J. Korean Earth Sci. Soc.* 42, 278–295. <https://doi.org/10.5467/jkess.2021.42.3.278>.
- Kumar, M., Dhangar, K., Thakur, A.K., Ram, B., Chaminda, T., Sharma, P., Kumar, A., Raval, N., Srivastava, V., Rinklebe, J., Kuroda, K., Sonne, C., Barcelo, D., 2021. Antidrug resistance in the Indian ambient waters of Ahmedabad during the COVID-19 pandemic. *J. Hazard Mater.* 416, 126125. <https://doi.org/10.1016/j.jhazmat.2021.126125>.
- Kumari, M., Kumar, A., 2020. Identification of component-based approach for prediction of joint chemical mixture toxicity risk assessment with respect to human health: a critical review. *Food Chem. Toxicol.* 143, 111458. <https://doi.org/10.1016/j.fct.2020.111458>.
- Kwok, C.S., Dashti, M., Tafuro, J., Nasiri, M., Muntean, E.-A., Wong, N., Kemp, T., Hills, G., Mallen, C.D., 2021. Methods to disinfect and decontaminate SARS-CoV-2: a systematic review of *in vitro* studies. *Therap. Adv. Inf. Dis.* 8 <https://doi.org/10.1177/2049936121998548>, 10.1177/2049936121998548.
- Li, Y., Zhou, C., Wang, S., Lin, Q., Ni, Z., Qiu, H., Morel, J.L., Qiu, R., 2019. Phytotoxicity and oxidative effects of typical quaternary ammonium compounds on wheat (*Triticum aestivum* L.) seedlings. *Environ. Sci. Pollut. Res. Int.* 26, 25985–25999. <https://doi.org/10.1007/s11356-019-05822-7>.
- Liu, Y.C., Kuo, R.L., Shih, S.R., 2020. COVID-19: the first documented coronavirus pandemic in history. *Biomed. J.* 43, 328–333. <https://doi.org/10.1016/j.bj.2020.04.007>.
- Ludovico, P., Burhans, W.C., 2014. Reactive oxygen species, ageing and the hormesis police. *FEMS Yeast Res.* 14, 33–39. <https://doi.org/10.1111/1567-1364.12070>.
- Marinello, S., Butturi, M.A., Gamberini, R., 2021. How changes in human activities during the lockdown impacted air quality parameters: a review. *Environ. Prog. Sustain. Energy* 40, e13672. <https://doi.org/10.1002/ep.13672>.
- Mo, L.Y., Liu, Y.A., Zhu, J., Qin, L.T., Liang, Y.P., Zeng, H.H., 2020. Benefits from hazards, benefits from nothing, and benefits from benefits: the combined effects of five quaternary ammonium compounds to *Vibrio qinghaiensis* Q67. *Environ. Sci. Eur.* 32, 35. <https://doi.org/10.1186/s12302-020-00310-z>.
- Nabi, G., Wang, Y., Hao, Y., Khan, S., Wu, Y., Li, D., 2020. Massive use of disinfectants against COVID-19 poses potential risks to urban wildlife. *Environ. Res.* 188, 109916. <https://doi.org/10.1016/j.envres.2020.109916>.
- Neagu, M., Calina, D., Docea, A.O., Constantin, C., Filippini, T., Vinceti, M., Drakoulis, N., Poulas, K., Nikolouzakis, T.K., Spandidos, D.A., Tsatsakis, A., 2021. Back to basics in COVID-19: antigens and antibodies—completing the puzzle. *J. Cell Mol. Med.* 25, 4523–4533. <https://doi.org/10.1111/jcmm.16462>.
- Pereira, B.M.P., Tagkopoulos, I., 2019. Benzalkonium chlorides: uses, regulatory status, and microbial resistance. *Appl. Environ. Microbiol.* 85, e00377-19 <https://doi.org/10.1128/aem.00377-19>.
- Pérez-Torrado, R., Querol, A., 2016. Opportunistic strains of *Saccharomyces cerevisiae*: a potential risk sold in food products. *Front. Microbiol.* 6, 1522. <https://doi.org/10.3389/fmicb.2015.01522>.
- Pichel, N., Vivar, M., Fuentes, M., 2019. The problem of drinking water access: a review of disinfection technologies with an emphasis on solar treatment methods. *Chemosphere* 218, 1014–1030. <https://doi.org/10.1016/j.chemosphere.2018.11.205>.
- Rabenau, H.F., Kampf, G., Cinatl, J., Doerr, H.W., 2005. Efficacy of various disinfectants against SARS coronavirus. *J. Hosp. Infect.* 61, 107–111. <https://doi.org/10.1016/j.jhin.2004.12.023>.
- Rodríguez-Mozaz, S., Chamorro, S., Martí, E., Huerta, B., Gros, M., Sánchez-Melsió, A., Borrego, C.M., Barceló, D., Balcázar, J.L., 2015. Occurrence of antibiotics and antibiotic resistance genes in hospital and urban wastewaters and their impact on the receiving river. *Water Res.* 69, 234–242. <https://doi.org/10.1016/j.watres.2014.11.021>.
- Schwarz, S., Loeffler, A., Kadlec, K., 2017. Bacterial resistance to antimicrobial agents and its impact on veterinary and human medicine. *Vet. Dermatol.* 28 <https://doi.org/10.1111/vde.12362>, 82–e19.
- Sedlak, D.L., Von Gunten, U., 2011. The chlorine dilemma. *Science* 331, 42–43. <https://doi.org/10.1126/science.1196397>.
- Semchyshyn, H.M., 2014. Hormetic concentrations of hydrogen peroxide but not ethanol induce cross-adaptation to different stresses in budding yeast. *Internet J. Microbiol.* 2014, 485792 <https://doi.org/10.1155/2014/485792>.
- Semchyshyn, H.M., Valishkevych, B.V., 2016. Hormetic effect of H₂O₂ in *Saccharomyces cerevisiae*: involvement of TOR and glutathione reductase. *Dose-Response* 14. <https://doi.org/10.1177/1559325816636130>.
- Shahid, M., Niazi, N.K., Rinklebe, J., Bundschuh, J., Dumat, C., Pinelli, E., 2020. Trace elements-induced phytohormesis: a critical review and mechanistic interpretation. *Crit. Rev. Environ. Sci. Technol.* 50, 1984. <https://doi.org/10.1080/10643389.2019.1689061>, 2015.
- Sharafi, S.M., Ebrahimpour, K., Nafez, A., 2021. Environmental disinfection against COVID-19 in different areas of health care facilities: a review. *Rev. Environ. Health* 36, 193–198. <https://doi.org/10.1515/reveh-2020-0075>.
- Shimabukuro, P.M.S., Duarte, M.L., Imoto, A.M., Atallah, Á.N., Franco, E.S.B., Peccin, M. S., Tamaino, M., 2020. Environmental cleaning to prevent COVID-19 infection. A rapid systematic review. *Sao Paulo Med. J.* 138, 505–514. <https://doi.org/10.1590/1516-3180.2020.0417.09092020>.
- Subpiramanyam, S., 2021. Outdoor disinfectant sprays for the prevention of COVID-19: are they safe for the environment? *Sci. Total Environ.* 759, 144289. <https://doi.org/10.1016/j.scitotenv.2020.144289>.
- Sun, T., Zhan, J., Li, F., Ji, C., W, H., 2021. Effect of microplastics on aquatic biota: a hormetic perspective. *Environ. Pollut.* 285, 117206. <https://doi.org/10.1016/j.envpol.2021.117206>.
- Tezel, U., Pavlostathis, S.G., 2011. Role of quaternary ammonium compounds on antimicrobial resistance in the environment. In: *Antimicrobial Resistance in the Environment*. Wiley-Blackwell, pp. 349–387. <https://doi.org/10.1002/9781118156247.ch20>.
- Tsatsakis, A., 2021. Toxicological Risk Assessment and Multi-System Health Impacts from Exposure, first ed. Academic Press. <https://doi.org/10.1016/c2020-0-02454-0>.
- Tsatsakis, A.M., Docea, A.O., Calina, D., Buga, A.M., Zlatian, O., Gutnikov, S., Kostoff, R. N., Aschner, M., 2019. Hormetic Neurobehavioral effects of low dose toxic chemical mixtures in real-life risk simulation (RLRS) in rats. *Food Chem. Toxicol.* 125, 141–149. <https://doi.org/10.1016/j.fct.2018.12.043>.
- Tsitsifli, S., Kanakoudis, V., 2018. Disinfection impacts to drinking water safety—a review. *SAVE Proc.* 2, 603. <https://doi.org/10.3390/proceedings2110603>.
- World Health Organization (WHO), 2020. *Cleaning and Disinfection of Environmental Surfaces in the Context of COVID-19: Interim Guidance*, 15 May 2020.
- Xiling, G., Yin, C., Ling, W., Xiaosong, W., Jingjing, F., Fang, L., Xiaoyan, Z., Yiyue, G., Ying, C., Lumbiao, C., Liubo, Z., Hong, S., Yan, X., 2021. In vitro inactivation of SARS-CoV-2 by commonly used disinfection products and methods. *Sci. Rep.* 11, 2418. <https://doi.org/10.1038/s41598-021-82148-w>.
- Zaman, S. Bin, Hussain, M.A., Nye, R., Mehta, V., Mamun, K.T., Hossain, N., 2017. A review on antibiotic resistance: alarm bells are ringing. *Cureus* 9, e1403. <https://doi.org/10.7759/cureus.1403>.
- Zhang, C., Cui, F., Zeng, G. ming, Jiang, M., Yang, Z., zhu, Yu, gang, Z., Zhu, M., ying, Shen, qing, L., 2015. Quaternary ammonium compounds (QACs): a review on occurrence, fate and toxicity in the environment. *Sci. Total Environ.* 518–519, 352–362. <https://doi.org/10.1016/j.scitotenv.2015.03.007>.
- Zhang, H., Tang, W., Chen, Y., Yin, W., 2020. Disinfection threatens aquatic ecosystems. *Science* 368, 146–147. <https://doi.org/10.1126/science.abb8905>.
- Zhang, Q., Pi, J., Woods, C.G., Jarabek, A.M., Clewell, H.J., Andersen, M.E., Andersen, M. E., 2008. Hormesis and adaptive cellular control systems. *Dose. Response* 6, 196–208. <https://doi.org/10.2203/dose-response.07-028.Zhang>.